



CEOP Model Output : MOLTS MSC data

1. IDENTIFICATION INFORMATION

Name	CEOP Model Output : MOLTS MSC data
Abbreviation	Coordinated Energy and Water-Cycle Observation Project Model Output (MOLTS) UKMO: MSC: Meteorological Service Canada data
Metadata Identifier	CEOP_Model_MOLTS_MSC20200901155745-DIAS20200901154929-en

2. CONTACT

2.1 CONTACT on DATASET

Name	Stephane Belair
Organization	Meteorological Service Canada
Address	77 Westmorland Street, Fredericton, New Brunswick, Suite 260, Canada
TEL	506-451-6010
FAX	819-994-0736

2.2 CONTACT on PROJECT

2.2.1 Data Integration and Analysis System

Name	DIAS Office
Organization	Remote Sensing Technology Center of Japan
Address	TOKYU REIT Toranomom Building 2F 3-17-1 Toranomom, Minato-ku, Tokyo, 105-0001, Japan
E-mail	dias-office@diasjp.net

3. DOCUMENT AUTHOR

Name	Stephane Belair
Organization	Meteorological Service Canada

4. DATASET CREATOR

Name	Stephane Belair
Organization	Meteorological Service Canada

5. DATE OF THIS DOCUMENT

2020-09-01

6. DATE OF DATASET

publication : 2010-04-10

7. DATASET OVERVIEW

7.1 Abstract

Ten operational Numerical Weather Prediction (NWP) and two data assimilation centers are currently contributing analysis/assimilation and forecast model products from global and regional NWP suites, including both operational and reanalysis systems to this component of CEOP. The contributing centers include:

BoM: Bureau of Meteorology

CPTEC: Centro de Previsao de Tempo e Estudos Climaticos

ECMWF: European Centre for Medium-Range Weather Forecasts

ECPC: Experimental Climate Prediction Center

EMC: EPSON Meteo Center (Centro EPSON Meteo)

GLDAS: Global Land Data Assimilation System

GMAO: NASA Global Modeling and Assimilation Office

JMA: Japan Meteorological Agency

MSC: Meteorological Service Canada

NCEP: National Centers for Environmental Prediction

NCMRWF: National Center for Medium Range Weather Forecasting

UKMO: UK Met Office

The Max-Planck Institute for Meteorology (MPIM) in coordination with the ICSU World Data Center for Climate (WDCC) in Hamburg, Germany was designated as the CEOP model output archive center. The WDCC is administered by the Model and Data Group (M&D) at MPIM and the German Climate Computing Center (DKRZ).

To assist with the organization of this activity during the Coordinated Enhanced Observing Period ('CEOP'), a Model Output Management Document was drafted as a guide for the participating centers to use in setting up their processes for meeting their commitments to 'CEOP'. The Guidance Document addressed the two issues of (1) the model output variables requested by 'CEOP' and (2) the two types of requested model output, namely global gridded (in GRIB format) and site-specific Model Output Location Time Series (MOLTS) at each of the 'CEOP' Reference Sites.

A new version of the Guidance Document will be compiled that clarifies what model output data will be generated by the NWP Centers and Groups contributing to the model output component of Coordinated Energy and Water Cycle Observations Project (CEOP) and how they will interface/transfer the data that will be handled and retained at the WDCC. The issues covered in the document will include: (1) global versus regional products; (2) desired assimilation output; Interval and length of free-

running forecasts; (3) Operational versus reanalysis data; (4) the CEOP schedule/archive periods; (5) the number and locations of MOLTS sites; and (6) the homogenizing of the model output and metadata formats (i.e. standard parameters).

Results up to this point in the CEOP model output generation effort make it clear that the transfer aspect of the data handling effort has been progressing well. Data from all twelve Centers participating in CEOP have been received at the data archive center and has either been placed into the database at the Hamburg facility, or is in the process of being entered into the database. The current data holdings in the MPIM archive can be viewed http://www.mad.zmaw.de/fileadmin/extern/wdc/ceop/Data_timeline_L_12.pdf.

7.2 Topic Category(IS019139)

climatologyMeteorologyAtmosphere

7.3 Temporal Extent

Begin Date	01-OCT-2002
End Date	31-DEC-2004
Temporal Characteristics	900sec

7.4 Geographic Bounding Box

North latitude bound	71.62
West longitude bound	148.15
Eastbound longitude	-156.62
South latitude bound	-35.66

7.5 Grid

7.6 Geographic Description

7.7 Keywords

7.7.1 Keywords on Dataset

Keyword Type	Keyword	Keyword thesaurus Name
theme	Climate	GEOSS

7.7.2 Keywords on Project

7.7.2.1 Data Integration and Analysis System

Keyword Type	Keyword	Keyword thesaurus Name
theme	DIAS > Data Integration and Analysis System	No_Dictionary

7.8 Online Resource

MSC : http://www.eol.ucar.edu/projects/ceop/dm/model/model_chars.html#msc

CEOP Distributed Data Integration System : http://jaxa.ceos.org/wtf_ceop/

CEOP Centralized Data Integration System : http://monsoon.t.u-tokyo.ac.jp/ceop-dc/ceop-dc_top.htm

GCMD CEOP Portal : <http://gcmd.gsfc.nasa.gov/KeywordSearch/Home.do?Portal=ceop&MetadataType=0>

CEOP Model Data Gateway : <http://www.mad.zmaw.de/projects-at-md/ceop/>

file download : <https://data.diasjp.net/dl/storages/filelist/dataset:27>

7.9 Data Environmental Information

7.10 Distribution Information

name	version	specification
------	---------	---------------

8. DATA PROCESSING

9. DATA REMARKS

For the atmospheric model GEM that was used for CEOP

Côté J., S. Gravel, A. Mailhot, A. Patoine, M. Roch, and A. Staniforth, 1998a: The operational CMC-MRB global environmental multiscale (GEM) model. Part I: Design considerations and formulation. Mon. Wea. Rev., 126, 1373-1393.

For the particular configuration of the global system that was used for CEOP:

Blaizot, S., J. Mailhot, C. Girard, and P. Vaillancourt, 2005: Boundary layer and shallow cumulus clouds in a medium-range forecast of large-scale weather system. Mon. Wea. Rev., 133, 1938-1960.

Blaizot, S., M. Roch, A.-M. Leduc, P.A. Vaillancourt, S. Laroche, and J. Mailhot, 2008: Medium-range quantitative precipitation forecasts from Canada's new 33-km deterministic global operational system. Wea. Forecasting (conditionally accepted, but the revisions required are small).

10. LICENSE

10.1 Data Policy by the Data Provider

10.2 Data Policy by the Project

10.2.1 Data Integration and Analysis System

If data provider does not have data policy, DIAS Terms of Service (<https://diasjp.net/en/policy/>) and DIAS Privacy Policy (<https://diasjp.net/en/privacypolicy/>) apply.

If there is a conflict between DIAS Terms of Service and data provider's policy, the data provider's policy shall prevail.

11. DATA SOURCE ACKNOWLEDGEMENT

11.1 Acknowledge the Data Provider

11.2 Acknowledge the Project

11.2.1 Data Integration and Analysis System

If you plan to use this dataset for a conference presentation, paper, journal article, or report etc., please include acknowledgments referred to following examples. If the data provider describes examples of acknowledgments, include them as well.

"We used the [name of dataset] provided by [name of data provider] in this study. This dataset was collected and provided under the Data Integration and Analysis System (DIAS, Project No. JPMXD0716808999), which has been developed and operated by the Ministry of Education, Culture, Sports, Science and Technology (MEXT)."

12. DISCLAIMER

12.1 Disclaimer of Project

12.1.1 Data Integration and Analysis System

If data provider does not have data policy, disclaimer of DIAS Terms of Service (<https://diasjp.net/en/policy/>) apply.

If there is a conflict between DIAS Terms of Service and data provider's policy, the data provider's policy shall prevail.

13. REFERENCES

For the atmospheric model GEM that was used for CEOP

Côté J., S. Gravel, A. M. Gauthier, A. Patoine, M. Roch, and A. Staniforth, 1998a: The operational CMC-MRB global environmental multiscale (GEM) model. Part I: Design considerations and formulation. *Mon. Wea. Rev.*, 126, 1373-1393.

For the particular configuration of the global system that was used for CEOP:

B lair, S., J. Mailhot, C. Girard, and P. Vaillancourt, 2005: Boundary layer and shallow cumulus clouds in a medium-range forecast of large-scale weather system. *Mon. Wea. Rev.*, 133, 1938–1960.

B lair, S., M. Roch, A.-M. Leduc, P.A. Vaillancourt, S. Laroche, and J. Mailhot, 2008: Medium-range quantitative precipitation forecasts from Canada’s new 33-km deterministic global operational system. *Wea. Forecasting* (conditionnally accepted, but the revisions required are small).

Copyright(c) 2006-2020 Data Integration & Analysis System (DIAS) All Rights Reserved. This project is supported by ” Data Integration & Analysis System ” funded by MEXT, Japan
--